



Treatment

Treatment aims to relieve pain, treat acute flare-ups by controlling inflammation, and prevent recurrence.

The 2025 French national guidelines propose:

▶ First-line treatment

Non-steroidal anti-inflammatory drugs (NSAIDs) (aspirin or ibuprofen) combined with colchicine.

Colchicine will be prescribed for 3 months in the case of a first episode and for at least 6 months in the case of recurrence.

▶ Second-line treatment

• Interleukin-inhibiting biotherapy (anakinra) may be offered by a specialist center for refractory forms only

▶ Treatment should also target the underlying cause if it is identified.

▶ Corticosteroid therapy is no longer recommended except in specific cases (lupus, post-surgery).

▶ Drainage is exceptional, reserved for abundant effusions with repercussions.



Our team:

Internal medicine

Prof. Sophie GEORGIN-LAVIALLE
Dr. Léa SAVEY
Dr. Marion DELPLANQUE
Dr. Catherine GRANDPEIX-GUYODO
Dr. Yves-Jean Zhu
Dr. Philippe MERTZ
Dr. Robin ECHERBAULT

Dermatology & Allergology

Prof. Angèle SORIA
Dr. Jean-Benoît MONFORT

Nephrology

Prof. Jean-Jacques BOFFA

Anatomical Pathology

Prof. David BUOB

Hepatogastroenterology

Dr. Ingrid POPA

Tenon Hospital
Internal Medicine Department

4 rue de la Chine
75020 Paris, France
Tel: +33 (0)1 56 01 74 31
Fax: 01 56 01 71 46


✉ ceremaia-medecine-int.tenon@aphp.fr

📷 @CEREMAIA_Tenon

<https://www.maladiesautoinflammatoires.fr/>



Recurrent pericarditis



<https://www.maladiesautoinflammatoires.fr/>

📷 @CEREMAIA_Tenon



Recurrent pericarditis

Definition

Recurrent pericarditis is the reappearance of acute pericarditis (inflammation of the membrane surrounding the heart) after an asymptomatic period of at least 4 to 6 weeks.

It can be caused by various factors, such as infections, particularly viral infections, immunoinflammatory diseases, or post-surgical conditions, but in most cases the cause is not identified, and it is then referred to as idiopathic and/or viral pericarditis.

Epidemiology

This is a rare disease. The risk of recurrence after acute pericarditis is estimated at 20 to 30%, and between 5 and 10% of patients are at risk of multiple recurrences.

It most often affects young or middle-aged individuals, with no gender predominance.



Etiology

The main causes of recurrent pericarditis are:

1. Infections:
 - Viral: such as coxsackivirus infections, herpes, HIV, or other viruses.
 - Bacterial: particularly tuberculosis.
 - Fungal or parasitic: rarer and mainly in immunocompromised individuals.
2. Diseases:
 - Autoimmune: such as systemic lupus erythematosus, rheumatoid arthritis, systemic scleroderma, or Sjögren's syndrome.
 - Autoinflammatory diseases such as familial Mediterranean fever or Still's disease, for example.
3. Chest trauma or chest or heart surgery can be complicated by acute pericarditis, which can develop into recurrent episodes.
4. Certain cancers such as lung cancer, breast cancer, or lymphomas.
5. When no cause is found, pericarditis is considered idiopathic (***the most common case***).



Clinical

Patients suffer from acute chest pain behind the sternum, which is worse when breathing deeply, with fever and sometimes a feeling of chest tightness and shortness of breath.



Diagnosis

The diagnosis of recurrent pericarditis is based on:

- The presence of pain. Typical chest pain +/- fever
- The presence of blood inflammation revealed by blood tests with **elevated** inflammation markers, mainly **C-reactive protein (CRP)**.
- Evidence of pericardial effusion (increased pericardial fluid) on cardiac ultrasound or chest CT scan or cardiac magnetic resonance imaging (MRI) in cases of diagnostic uncertainty.

In rare cases, pericardial fluid puncture or even pericardial biopsy may be performed to confirm the diagnosis and investigate specific causes, particularly if inflammatory or infectious diseases are suspected.

If CRP is not elevated, reconsider the diagnosis and seek expert advice.

